

DIAMMONIUM PHOSPHATE

Crop and/or situation (a)	F G I (b)	Pests or group of pests controlled (c)	Formulation		Application				Application rate per treatment			Total rate	PHI (days) (m)	Remarks				
			Type (d-f)	Conc of a.s. g/L (i)	Method kind (f-h)	Growth stage and season (j)	Number min max (k)	Interval between applications (min)	kg a.s./hl min max (kg/hl)	Water l/ha min max	kg a.s./ha min max (kg/ha) (l)				kg a.s./ha min max (kg/ha) (l)			
Orchards Including cherry tree <i>Prunus spp</i>	F	Mediterranean Fruit fly <i>Ceratitis capitata</i> Cherry fly <i>Rhagoletis cerasi</i>	VP	40	Placed in physical traps		Mass trapping: 1 trap per tree up to 100 traps/ha	Approx. 6 to 8 Weeks*	max 4	Mass trapping: max 100	Mass trapping: max 4	Mass trapping: max 4	Not relevant					
Olive trees <i>Olea europaea</i>		Olive fly <i>Bactrocera oleae</i>													Mass trapping: 1 trap per tree up to 100 traps/ha	Mass trapping: max 100	Mass trapping: max 4	Mass trapping: max 4
<i>Citrus spp</i>		Mediterranean Fruit fly <i>Ceratitis capitata</i>													Mass trapping: 1 trap per tree up to 100 traps/ha	Mass trapping: max 100	Mass trapping: max 4	Mass trapping: max 4
Other crops where <i>C. capitata</i> cause damage		Mediterranean Fruit fly <i>Ceratitis capitata</i>													Mass trapping: 1 trap per tree up to 100 traps/ha	Mass trapping: max 100	Mass trapping: max 4	Mass trapping: max 4

* depending upon environmental factors such as climate and topography

- (a) For crops, the EU and Codex classification (both) should be taken into account ; where relevant, the use situation should be described (e.g. fumigation of a structure)
- (b) Outdoor or field use (F), greenhouse application (G) or indoor application (I)
- (c) e.g. pests as biting and sucking insects, soil born insects, foliar fungi, weeds or plant elicitor
- (d) e.g. wettable powder (WP), emulsifiable concentrate (EC), granule (GR) etc..
- (e) GCPF Codes – GIFAP Technical Monograph N° 2, 1989
- (f) All abbreviations used must be explained
- (g) Method, e.g. high volume spraying, low volume spraying, spreading, dusting, drench
- (h) Kind, e.g. overall, broadcast, aerial spraying, row, individual plant, between the plant – type of equipment used must be indicated
- (i) g/kg or g/L. Normally the rate should be given for the active substance (according to ISO)
- (j) Growth stage at last treatment (BBCH Monograph, Growth Stages of Plants, 1997, Blackwell, ISBN 3-8263-3152-4), including where relevant, information on season at time of application
- (k) Indicate the minimum and maximum number of application possible under practical conditions of use
- (l) The values should be given in g or kg whatever gives the more manageable number (e.g. 200 kg/ha instead of 200 000 g/ha or 12.5 g/ha instead of 0.0125 kg/ha)
- (m) PHI - minimum pre-harvest interval