

SALIX SPP CORTEX

Crop and/or situation (a)	Example product name as available on the market	F G I (b)	Target (c)	Product		Application				Application rate per treatment			Total rate	PHI (days) (m)	Remarks (*)
				Type (d-f)	Conc of a.i. g/kg (i)	Method kind (f-h)	Growth stage and season** (j)	Number min max (k)	Interval between applications (min)	g a.i./hl min max (g/hl)	Water l/ha min max	g a.i./ha min max (g/ha) (l)			
Fruit trees Peach-tree <i>Prunus persica</i>	Homogenate of <i>Salix</i> spp	F	Foliar fungi like <i>Taphrina deformans</i>	Dispersible concentrate (DC)	2.22	Foliar application spraying	From 1 <sup>st</sup> Shoots (BCH10) to cluster tightening (BBCH57) Spring	2 to 6	7 days	222.22	500 to 1000	1111.11 to 2222.22	2.22 to 13.33	None	Plant Homogenate extracted with hot water (infusion), filtered and diluted by 3, to be used up to a maximum of 24 h after preparation The product cannot be applied in case of hot temperature. It is used in case of rainy period;
Apple fruit <i>Malus pumila</i> , <i>Malus domestica</i>		F	Foliar fungi like scab disease <i>Venturia inaequalis</i> Powdery mildews: <i>Podosphaera leucotricha</i>				From green leaf tip (BBCH 53) to flowers fading (BBCH 67) Spring								
Grapevine <i>Vitis vinifera</i>		F	Downy mildews: <i>Plasmopara viticola</i> , Powdery mildews: <i>Erysiphe necator</i>				From 1 <sup>st</sup> shoots (BCH10) to cluster tightening (BBCH57) Spring to summer								

\*\*\* \*\*

\* For uses where the column „Remarks. As above or other conditions to take into account

(a) For crops, the EU and Codex classification (both) should be taken into account ; where relevant, the use situation should be described (e.g. fumigation of a structure)

(b) Outdoor or field use (F), greenhouse application (G) or indoor application (I)

(c) e.g. pests as biting and sucking insects, soil born insects, foliar fungi, weeds or plant elicitor

(d) e.g. wettable powder (WP), emulsifiable concentrate (EC), granule (GR) etc...

(e) GCPF Codes – GIFAP Technical Monograph N° 2, 1989

(f) All abbreviations used must be explained

(g) Method, e.g. high volume spraying, low volume spraying, spreading, dusting, drench

(h) Kind, e.g. overall, broadcast, aerial spraying, row, individual plant, between the plant – type of equipment used must be indicated

(i) g/kg or g/L. Normally the rate should be given for the active substance (according to ISO)

(j) Growth stage at last treatment (BBCH Monograph, Growth Stages of Plants, 1997, Blackwell, ISBN 3-8263-3152-4), including where relevant, information on season at time of application

(k) Indicate the minimum and maximum number of application possible under practical conditions of use

(l) The values should be given in g or kg whatever gives the more manageable number (e.g. 200 kg/ha instead of 200 000 g/ha or 12.5 g/ha instead of 0.0125 kg/ha)

(m) PHI - minimum pre-harvest interval