

SUCROSE

Crop and/or situation (a)	Member State	Example product name as available on the market	F G I (b)	Target (c)	Product		Application				Application rate per treatment			Total rate g a.i./ha min max (g/ha) (l)	PHI (days) (m)	Remarks (*)
					Type (d-f)	Conc of a.i. g/kg (i)	Method kind (f-h)	Growth stage and season** (j)	Number min max (k)	Interval between applications (min)	g a.i./hl min max (g/hl)	Water l/ha min max	g a.i./ha min max (g/ha) (l)			
<i>Apple fruit Malus pumila Malus domestica</i>	France	Sucrose	F	fruits borer like Codling moth: <i>Cydia pomonella</i>	Water Soluble Powder (SP)	998 to 1000	Foliar application spraying early in the morning before 9 AM (solar time)*	From spring BBCH stage 6 to Summer stage 65	7 to 10	15 days*	1	600 to 1000	6 to 10	42 to 100	None	Cold Water Solution prepared just before application
<i>Sweet Maize (Sweet corn) Zea mays</i>			F	Corn borer: <i>Ostrinia nubilalis</i> Hbn.,			From the BBCH stage 12 to 51	3 to 4	15 days*	200		2	6 to 8			

- * For uses where the column „Remarks. As above or other conditions to take into account
- (a) For crops, the EU and Codex classification (both) should be taken into account ; where relevant, the use situation should be described (e.g. fumigation of a structure)
- (b) Outdoor or field use (F), greenhouse application (G) or indoor application (I)
- (c) e.g. pests as biting and sucking insects, soil born insects, foliar fungi, weeds or plant elicitor
- (d) e.g. wettable powder (WP), emulsifiable concentrate (EC), granule (GR) etc..
- (e) GCPF Codes – GIFAP Technical Monograph N° 2, 1989
- (f) All abbreviations used must be explained
- (g) Method, e.g. high volume spraying, low volume spraying, spreading, dusting, drench
- (h) Kind, e.g. overall, broadcast, aerial spraying, row, individual plant, between the plant – type of equipment used must be indicated
- (i) g/kg or g/L. Normally the rate should be given for the active substance (according to ISO)
- (j) Growth stage at last treatment (BBCH Monograph, Growth Stages of Plants, 1997, Blackwell, ISBN 3-8263-3152-4), including where relevant, information on season at time of application
- (k) Indicate the minimum and maximum number of application possible under practical conditions of use
- (l) The values should be given in g or kg whatever gives the more manageable number (e.g. 200 kg/ha instead of 200 000 g/ha or 12.5 g/ha instead of 0.0125 kg/ha)
- (m) PHI - minimum pre-harvest interval