

WHEY

Crop and/or situation (a)	F G or I (b)	Pests or group of pests controlled (c)	Formulation		Application				Application rate			PHI (days)	Remarks
			Type (d-f)	Conc. of a.i. g/L (i)	Method kind (f-h)	Growth stage & season (j)	No. of application min/max (k)	Interval between applications (min)	g a.i./hl min max (g/hl)	Water l/ha min max	Total rate each application g a.i./ha min max (g/ha) (l) or concentration recommended		
Cucumber <i>Cucumis Sativus</i> Zucchini squash <i>Cucurbita pepo</i>	G	Powdery mildews: <i>Podosphaera fusca</i> <i>Podosphaera xanthii</i> <i>Golovinomyces/ Erysiphe cichoracearum</i> and <i>orontii</i> <i>Sphaerotheca fuliginea</i> <i>Leveillula cucurbitacearum</i>	TC	60-80 g/L	Foliar spray*	From three weeks after sowing (9th leaf unfolded on main stem) to 9 or more primary side shoots visible (BBCH 19-49)**	3 - 5	7 days	0.6 L to 3 L (0.036 to 0.24 kg a.s.)	1000 to 1500	6 to 30 L (0.36 to 2.4 kg a.s.)		Whey should be used rapidly after collection, not stored in metal vessel.

* spray when there is sun (preferably morning)

** do not apply when any plant in the greenhouse is at a later growth stage than BBCH 49.

(a) For crops, the EU and Codex classification (both) should be taken into account ; where relevant, the use situation should be described (e.g. fumigation of a structure)	(i) g/kg or g/L. Normally the rate should be given for the substance (according to ISO)
(b) Outdoor or field use (F), greenhouse application (G) or indoor application (I)	(j) Growth stage at last treatment (BBCH Monograph, Growth Stages of Plants, 1997, Blackwell, ISBN 3-8263-3152-4), including where relevant, information on season at time of application
(c) e.g. pests as biting and sucking insects, soil born insects, foliar fungi, weeds or plant elicitor	(k) Indicate the minimum and maximum number of application possible under practical conditions of use
(d) e.g. wettable powder (WP), emulsifiable concentrate (EC), granule (GR) etc..	(l) The values should be given in g or kg whatever gives the more manageable number (e.g. 200 kg/ha instead of 200 000 g/ha or 12.5 g/ha instead of 0.0125 kg/ha)
(e) GCPF Codes – GIFAP Technical Monograph N° 2, 1989	(m) PHI - minimum pre-harvest interval
(f) All abbreviations used must be explained	
(g) Method, e.g. high volume spraying, low volume spraying, spreading, dusting, drench	
(h) Kind, e.g. overall, broadcast, aerial spraying, row, individual plant, between the plant – type of equipment used must be indicated	