Data were assembled with care but treat differences below LSD cautiously. Performance variation is not Crop Evaluation or HGCA's responsibility. The control is the mean of Biststar, Lagoon and Taurus.

SPRING LINSEED

Result from the Cambridge trial FINAL RESULTS									
	Gross output		Seed yield		Oil content	Standing abitilty	Plant height	Ealiness of flowering	Earliness of maturity
Variety	t/ha	% control	t/ha	% control	(%)	(1-9)	(cm)	(1-9)	(1-9)
Control (t/ha)		1.9		1.9	, ,	` ,			
Juliet	2.41	125	2.30	124	43.0	4.0	58	6.7	4.0
Bilton	1.94	101	1.88	101	41.6	9.0	55	5.0	5.0
Biltstar	2.09	108	2.00	108	42.3	6.0	53	9.0	5.0
Talon	1.95	101	1.92	104	39.6	9.0	54	7.0	5.0
Sunrise	1.79	93	1.73	93	41.6	8.0	61	7.0	8.0
Lagoon	1.89	98	1.80	98	42.7	8.0	49	8.3	6.3
Linus	1.74	90	1.68	91	42.0	9.0	56	7.0	6.0
Taurus	1.8	93	1.74	94	41.4	7.0	58	7.3	8.7
Bethune	1.63	84	1.57	85	41.2	8.0	60	6.0	7.7
Oscar	2.21	115	2.17	117	39.7	7.0	53	8.0	6.0
Marmalade	1.78	93	1.71	93	42.1	8.3	56	6.3	7.0
Laser	1.54	80	1.48	80	41.7	9.0	51	5.7	6.7
Gemini	1.63	85	1.58	85	41.5	9.0	55	4.3	5.0
Thirlmere	2.03	105	1.91	103	44.2	9.0	56	5.7	5.7
Windermere	1.89	98	1.79	97	43.5	8.0	56	5.0	6.0
LSD 5% 9.6				9.7					

The HGCA Recommended List is independently managed by Crop Evaluation Limited

The second linseed trial in 2004 was abandoned