#### **AHDB Recommended List (Table 13b)**

## Winter oilseed rape 2017/18 - North region

YIELD, QUALITY, AGRONO	MY A	ND D	ISEA	SE R	ESIS	TANC	Έ										<b>Described Variety</b>
RECOMMENDED				*						*	*		NEW		*		
AHDB	Anastasia	Alizze	Campus	DK Explicit	Nikita	Barbados	V316-OL ∼	SY Harnas	V-324-0L ~	DK Exentiel	PT234	Mentor \$	DK Secret #	Amalie ¥	Troy#	Average LSD (5%)	DK Imagis-CL &
Variety type	Conv	RH	Conv	RH	Conv	Conv	RH	RH	RH	RH	RH	RH	RH	Conv	RH		RH
Scope of recommendation	N	UK	UK	N	UK	N	UK	UK	N	N	N	Sp	UK	Sp	UK		N
Gross output, yield adjusted for oil co	ntent (%	% contr	ol)														
North region (5.3 t/ha)	110	109	109	109	109	109	109	108	107	106	104	100	97	96	95	8.1	102
East/West region (5.4 t/ha)	-	108	106	[101]	108	100	108	104	104	[101]	[101]	101	100	98	94	4.8	[99]
United Kingdom (5.4 t/ha)	-	108	107	102	109	101	108	105	104	101	101	101	100	98	94	4.5	99
Seed yield (% control)																	
North region (4.9 t/ha)	111	108	109	108	108	109	108	110	106	106	103	99	99	96	96	7.6	101
East/West region (5.0 t/ha)	-	107	106	[101]	107	100	108	106	103	[102]	[101]	100	101	99	96	4.5	[99]
United Kingdom (5.0 t/ha)	-	107	107	102	107	101	108	106	103	103	101	100	101	99	96	4.2	99
Agronomic features																	
Resistance to lodging (1-9)	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	0.2	8
Stem stiffness(1-9)	8	8	8	7	8	8	8	7	8	8	8	8	8	8	8	0.5	8
Shortness of stem (1-9)	7	6	6	5	7	6	6	7	6	6	6	7	9	7	8	0.2	6
Earliness of flowering (1-9)	6	8	6	6	7	5	7	7	6	7	7	6	7	6	6	0.4	5
Earliness of maturity (1-9)	5	5	5	5	5	4	5	5	5	6	6	5	6	5	5	0.4	5
Seed quality (at 9% moisture)																	
Oil content, fungicide-treated (%)	45.1	46.0	45.7	46.2	46.3	45.5	46.0	44.5	46.3	45.0	45.8	46.2	44.5	44.9	44.6	0.3	45.3
Glucosinolate (µmoles/g of seed)	11.1	13.0	11.2	10.4	9.5	12.0	12.9	12.3	9.9	11.4	10.4	10.2	10.7	13.7	12.0	-	11.7
Disease resistance																	
Light leaf spot (1-9)	6	7	6	6	7	7	6	6	7	6	6	6	7	6	6	0.7	6
Stem canker (1-9)	6	5	6	9	4	7	5	5	4	8	5	3	8	6	4	1.2	5
Varieties no longer listed in the North	rogioni	Crook		-41.10	d DT04	4											

Varieties no longer listed in the North region: Cracker, Incentive and PT211

C = yield control (for current table). For this table, PR46W21, DK Cabernet, PT211 and Trinity were the yield controls.

On the 1-9 scales, high figures indicate that a variety shows the character to a high degree (eg high resistance).

The target (spring) plant population is 40 plants/m² for RL trials. Maximum seed rate is 70 seeds/m² and may be lower if conditions permit. Glucosinolate contents are taken from the National List trials data.

UK = recommended for both the East/West and North regions

N = recommended for the North region

Sp = specific recommendation

Conv = conventional open-pollinated variety RH = restored hybrid \* = variety no longer in trial in region

[] = limited data

LSD = least significant difference

Average LSD (5%): Varieties that are more than one LSD apart are significantly different at the 95% confidence level.

<sup>~</sup> HOLL (high oleic, low linolenic) variety

<sup>¥ =</sup> Amalie has a specific recommendation for its resistance to Turnip Yellows Virus (TuYV)

<sup>\$</sup> Mentor is recommended for growing on land infected with common strains of clubroot; Mentor may, however, be infected by some strains and infections that have been reported in some fields.

<sup>#</sup> Semi-dwarf varieties believed to carry the Bzh dwarfing gene in the heterozygous state but this has not been verified in RL tests

<sup>&</sup>amp; Clearfield® variety, with tolerance to specific imidazolinone herbicides

### **AHDB Recommended List (Table 14)**

# Winter oilseed rape 2017/18 - North region

Supplementary data																	<b>Described Variety</b>
RECOMMENDED				*						*	*		NEW		*		
AHDB	Anastasia	Alizze	Campus	DK Explicit	Nikita	Barbados	V316-OL ~	SY Harnas	V-324-0L ∼	DK Exentiel	PT234	Mentor \$	DK Secret #	Amalie ¥	Troy #	Average LSD (5%)	DK Imagis-CL &
Variety type	Conv	RH	Conv	RH	Conv	Conv	RH	RH	RH	RH	RH	RH	RH	Conv	RH		RH
Scope of recommendation	N	UK	UK	N	UK	N	UK	UK	N	N	N	Sp	UK	Sp	UK		N
Breeder/ UK contact		O. C	Oit		0.1		OIT	Oit	- 1			ОР	O.C	Op	OI C		
Breeder	Lim	R2n	Mom	DK	LimEur	Mom	Mon	SyP		Mon	DP	NPZ	MonTec	LimEur	LSPB		DK
UK contact	Lim	RAGT	KWS	DK	Lim	KWS	Mon	Syn	Mon	DK	DP	LSPB	Mon	Lim	DSV		DK
Annual treated gross output, yield adju	usted fo	or oil co	ntent (	% cont	trol) - N	orth reg	gion										
2013 (5.2 t/ha)	[111]	[105]	[98]	[105]	[106]	[111]	[99]	[104]	[107]	[104]	[101]	[97]	-	[90]	[97]	12.5	[100]
2014 (5.4 t/ha)	[113]	[116]	[115]	[115]	[113]	[111]	[115]	[118]	[109]	[108]	[110]	[100]	[95]	-	96	12.1	[108]
2015 (6.0 t/ha)	[106]	[106]	[106]	[101]	[104]	[104]	[105]	[103]	[105]	[101]	[101]	[103]	[96]	[97]	94	9.3	[95]
2016 (4.6 t/ha)	[115]	[110]	[113]	[116]	[113]	[112]	[113]	[107]	[109]	[111]	[103]	[99]	[100]	[95]	[94]	14.5	[104]
Agronomy																	
Plant height (cm)	145	151	154	164	145	152	157	147	155	153	151	147	119	147	130	3.1	158
Status in RL system																	
Year first listed	13	16	15	15	16	16	15	15	16	15	15	15	17	16	13		-
RL status	-	P2	-	*	P2	P2	-	-	P2	*	*	-	P1	P2	*		-

[] = limited data

#### Key to Breeder and UK contact codes

DK = DEKALB (www.dekalb.co.uk)
DP = DuPont Pioneer (www.pioneer.com/uk)

DSV = DSV UK (www.dsv-uk.co.uk) KWS = KWS UK (www.kws-uk.com)

LimEur = Limagrain Europe SA (www.limagrain.co.uk)

P1 = first year of recommendation

Lim = Limagrain UK (www.limagrain.co.uk)

LSPB = LS Plant Breeding (www.lspb.eu)
Mom = Momont, France

wom = womont, France

Mon = Monsanto UK Ltd (www.monsanto.com)

MonTec = Monsanto Technology LLC (www.monsanto.com)

P2 = second year of recommendation

NPZ = NPZ-Lembke, Germany (www.npz.de)

RAGT = RAGT Seeds (www.ragt.co.uk)

R2n = RAGT, France (www.ragt.co.uk)

Syn = Syngenta UK Ltd (www.syngenta.co.uk)

SyP = Syngenta Participations AG (www.syngenta.co.uk)