



***Avoiding
lodging in
winter wheat
– practical guidelines***





February 2005


Background

How crops lodge

Cereals lodge due to interactions between **wind**, **rain** and **soil** with the plant.



Stem strength depends on stem diameter and the composition and width of the stem wall.



Anchorage depends on the spread and depth of the root plate and the strength of surrounding soil.



Stem lodging occurs when the stem base has insufficient strength to hold the shoot up against leverage.



Root lodging occurs when the root system has insufficient anchorage to hold the plant up against leverage.

Costs of lodging

On average, every three or four years widespread lodging occurs on up to 20% of the UK winter wheat area. Lodging has serious economic effects; yield is reduced and quality impaired, while combining and drying costs increase.


Economic loss is greatest when crops lodge early. Up to 75% of yield can be lost if crops lodge at ear emergence. Later lodging has less effect on yield, but will impair quality characters such as Hagberg falling number. Lodged crops often require extra drying.

Factors affecting lodging

These guidelines describe how lodging risk, or standing power, is altered by the environment and crop management. Lodging risk is scored on the 1–9 scale used in the *Recommended List* for winter wheat.


Environmental factors

Wind speed



Lodging risk increases as the wind leverage exerted on shoots rises. A 5kph increase in the maximum wind speed has the effect of reducing the varietal lodging resistance score by 1.

Rainfall



Just 6–11mm of rain in a day reduces soil surface strength and significantly increases the risk of root lodging. In a season with four extra days (compared with the farm's average) when rainfall exceeds 7mm, the varietal lodging resistance score will effectively be reduced by 1.

Soil strength

Soil shear strength has a large impact on anchorage strength and has most effect on root lodging risk. Soils with good crumb structure, low clay content or high organic matter tend to provide less anchorage and hence crops are at a higher lodging risk.

Soil strength also reduces significantly as moisture content increases.

Other factors

Long growing seasons, high fertility sites or eyespot infection can increase lodging risk. The varietal lodging resistance score decreases by 0.5 points for each 1t/ha yield increase above 9t/ha.

Crops grown in fields with light and drought-prone soils tend to lodge less.

Latitude influences sowing date and length of growing season which affects lodging risk (see table).

Predicting lodging

Crop condition in spring can indicate future lodging risk.

- n **Root lodging** risk increases when plant population exceeds 200 plants/m².
- n **Stem lodging** risk increases when the canopy at GS31 is large (green area index of 2 or more, or ground cover of >60%) as such crops tend to be weak-stemmed.

Assessing and managing risk... ..stem a

To assess the likely threat of lodging to a specific crop, start with the resistance to lodging score for the variety in

Factor	Impact
<p>Variety</p> <p>The Recommended List provides a combined root and stem lodging resistance rating for each variety. Research shows that husbandry affects root and stem lodging to different extents.</p>	<p>A variety's risk of lodging is given on a 1–9 score. Varieties with a score without PGR of 7 or less can be considered "at risk".</p>
<p>Soil residual nitrogen</p> <p>The amount of nitrogen left in the soil after a previous crop can significantly affect lodging risk. See the AHDB Nutrient Management Guide (RB209) for details on calculating crop N need and soil residual nitrogen.</p>	<p>High levels of soil N (SNS index 3 or above) promote thick, dense crop canopies that are more susceptible to stem lodging. Each unit increase on the soil nitrogen supply (SNS) index reduces stem lodging resistance score by 1.5. The effect on root lodging is about 0.75. Early-sown crops (before October) are affected most by high soil N levels.</p>
<p>Sowing date</p> <p>Early sowing leads to taller crops with a higher risk of both root and stem lodging. They also have denser canopies and often weaker stems. Adjustments (see right) are for central England. In northern Britain treat score for end September as zero.</p>	<p>For each week's delay in sowing, between mid-September and early November, a crop's lodging resistance score increases by 0.5.</p>
<p>Plant population</p> <p>Crops grown at higher plant populations have an increased risk of root lodging due to reduced anchorage strength. Stem strength is also reduced to some extent.</p>	<p>At plant populations of 200–400 plants/m², each increase of 50 plants/m² reduces stem lodging resistance score by 0.5 and root lodging by 1.0. Effects are greater below 200 plants/m².</p>

Use stem and root lodging totals to identify how you can avoid either type of lodging.

Spring management action

	Total lodging resistance score
	Lodging risk
<p>Fertiliser nitrogen</p> <p>Early (pre-GS31) or excessive applications of nitrogen increase tiller numbers and reduce stem strength. This increases the risk of both stem and root lodging. Nitrogen applications should always seek to optimise yield and quality.</p>	<p>A canopy management approach to nitrogen use (delayed and reduced applications) increases both root and stem lodging resistance by 0.5 points. Delaying the first split where soil fertility is moderate or high is unlikely to affect yield.</p>
<p>Plant growth regulators</p> <p>Tall crops are more prone to root and stem lodging. Plant growth regulators shorten crops to reduce both lodging risks to similar extents.</p>	<p>A single PGR application can increase both root and stem lodging resistance by 0.5 to 1.5, depending on variety and conditions.</p>
<p>Spring rolling</p> <p>Loosely structured soils provide weak anchorage for crops. Spring rolling, in appropriate conditions, consolidates such soils to reduce root lodging risk.</p>	<p>Spring rolling before GS30 increases the root lodging resistance score by 1. Rolling after GS30 can damage the growing point.</p>

End root lodging may need different actions

in the field and then adjust that score for the crop's soil residual nitrogen, sowing date and plant population.

Look up your lodging resistance score without PGR from the AHDB Recommended List.

Write that same score in both boxes 'A'.

1 2 3 4 5
Resistance score without PGR

6 7 8 9
Resistance score without PGR

STEM LODGING

ROOT LODGING

SNS Index
4 3 2 1 0
-3 -1.5 0 +1.5 +3
Adjustment for your SNS index

SNS Index
4 3 2 1 0
-1.5 -0.75 0 +0.75 +1.5
Adjustment for your SNS index

Mid Sept End Sept Mid Oct End Oct Nov onwards
-2 -1 0 +1 +2
Adjustment for your sowing date

Mid Sept End Sept Mid Oct End Oct Nov onwards
-2 -1 0 +1 +2
Adjustment for your sowing date

Plants/m²
>400 400-300 300-200 200-150 <150
-1.75 -1 0 +0.75 +1.5
Adjustment for your plant population

Plants/m²
>400 400-300 300-200 200-150 <150
-3.5 -2 0 +1.5 +3
Adjustment for your plant population

Total

Use this total in the table below for guidance about reducing lodging risk

Total

Use this total in the table below for guidance about reducing lodging risk

below 5	5-6.8	7-8.8	9-10	over 10
v. high	high	medium	low	very low
Delay & reduce N	Delay & reduce N	Delay N		
Full PGR	Full PGR	Single PGR	PGR if high yield forecast	

below 5	5-6.8	7-8.8	9-10	over 10
v. high	high	medium	low	very low
Reduce N	Reduce N			
Full PGR	Full PGR	Single PGR	PGR if high yield forecast	
Roll	Roll	Roll		

For very high risk situations, preventative techniques may not always be sufficient to prevent lodging. Alterations to variety and cropping strategy may be required.

Plant growth regulator products

PGRs are generally most effective in good growing conditions. Splitting applications increases the chance of matching application timing with appropriate growing conditions.

Consult your buyer or processor before applying any PGR.

Applying a sequence of a gibberellic acid inhibitor followed by an ethylene producer (see below) will usually have at least an additive effect. In some instances, sequences may have more than the additive effect of individual treatments.

Before use, always read the label. If in doubt consult a BASIS-registered advisor.

Acknowledgments

These guidelines were funded by AHDB and are based on R&D information provided by John Spink and Pete Berry, ADAS. AHDB is grateful to the following people for their comments on the guidelines: Hazel Doonan, AIC; Colin Mountfield-Smith, BASF; Dr Jim McVittie, AHDB and Peter Harding, Mandops.

Further information

AHDB Wheat growth guide: cereals.ahdb.org.uk/wheatgg

If you want more information about AHDB you can contact us in the following ways:

AHDB, Stoneleigh Park, Kenilworth, Warwickshire CV8 2TL

T: 024 7669 2051

E: comms@ahdb.org.uk

Twitter: @TheAHDB

ahdb.org.uk

While the Agriculture and Horticulture Development Board seeks to ensure that the information contained within this document is accurate at the time of printing, no warranty is given in respect thereof and, to the maximum extent permitted by law, the Agriculture and Horticulture Development Board accepts no liability for loss, damage or injury howsoever caused (including that caused by negligence) or suffered directly or indirectly in relation to information and opinions contained in or omitted from this document.

Reference herein to trade names and proprietary products without stating that they are protected does not imply that they may be regarded as unprotected and thus free for general use. No endorsement of named products is intended, nor is any criticism implied of other alternative but unnamed products.