





Final wheat quality diagrams

The Cereal Quality Diagrams below show the proportion of Group 1 and 2 samples meeting or exceeding the typical standards for high and medium quality bread wheat. [Use the Cereal Quality Calculator](#) to check your own specifications for this year's crop, where you can also [read an explanation of our Cereal Quality Diagrams](#).



In order to ensure a representative sample covering regions, varieties and data providers is given, some samples have been removed by stratified random selection. Please note, that due to the selection method, samples from a previous release may not be included within these results.

Further details on the AHDB Cereals and Oilseeds Cereal Quality Survey can be found at <http://cereals.ahdb.org.uk/markets/survey-results.aspx>

For more information, contact: Eleanor Holdsworth, Marcomms Executive, AHDB Cereals & Oilseeds, T: 024 7647 8753, E: Eleanor.Holdsworth@ahdb.org.uk

CEREAL QUALITY SURVEY RESULTS 2017

Wheat Analysis by Variety

	nabim group	Number of samples	Mean Values									
			Moisture Content		Specific Weight		Hagberg F-Number				Protein Content	
			% d.m	s.d*	Kg/hl	s.d*	Seconds	s.d*	Min	Max	% d.m	s.d*
Skyfall	1	4,986	14.9	1.01	76.8	2.06	248	76.01	91	401	13.2	0.89
Crusoe	1	4,481	14.8	0.98	75.8	2.14	245	72.85	91	401	13.7	0.75
Gallant	1	544	14.8	0.94	76.4	2.07	314	53.88	117	401	13.5	0.84
All Grp 1	1	11,426	14.9	1.00	76.3	2.16	254	75.25	91	401	13.4	0.87
Cordiale	2	1,359	14.8	0.97	77.2	1.96	304	59.05	98	401	13.4	0.79
KWS-Siskin	2	1,347	14.8	0.97	75.6	1.97	240	72.85	91	391	12.2	0.83
KWS-Lili	2	1,204	14.9	1.01	75.5	1.85	265	64.66	91	401	12.0	0.81
All Grp 2	2	3,949	14.8	0.98	76.1	2.08	271	70.61	91	401	12.5	1.02
Zulu	3	790	14.9	1.01	74.9	2.07	199	59.81	91	401	12.0	0.83
Claire	3	379	14.7	0.99	75.6	2.10	229	69.95	91	367	12.0	0.75
KWS-Basset	3	333	14.8	0.92	75.6	1.69	217	47.19	93	330	12.0	0.72
All Grp 3	3	1,848	14.9	1.00	75.2	1.98	212	59.41	91	401	12.0	0.79
JB-Diego	4	1,282	14.9	1.00	75.7	1.95	279	64.14	93	401	11.9	0.78
All Grp 4	4	5,466	14.9	0.99	75.3	2.23	231	71.54	91	401	11.9	0.80
ukp	1 or 2	9,213	14.8	0.98	76.0	2.12	260	72.90	91	401	13.2	1.05
uks	3 or 4	3,584	14.8	1.00	75.2	2.02	213	57.30	91	401	11.9	0.82

* Standard Deviation

All averages are simple arithmetic means
Results shown for 100+ samples only

Wheat Analysis by Region

Mean Values

	nabim group	Number of samples	Moisture Content		Specific Weight		Hagberg F-Number		Protein Content	
			% d.m	s.d*	Kg/hl	s.d*	Seconds	s.d*	% d.m	s.d*
South East	1	3,180	14.9	1.06	76.2	2.21	230	84.52	13.5	0.82
	2	516	14.9	0.99	76.4	2.20	246	81.89	12.8	1.12
	3	371	14.8	0.97	75.4	1.98	203	71.96	12.0	0.69
	4	534	14.7	1.10	75.4	2.24	228	75.10	12.0	0.76
	ukp	2,330	14.9	1.02	76.0	2.18	230	79.69	13.4	0.96
	uks	650	14.8	1.02	75.2	2.06	206	65.20	12.2	0.77
South West	1	1,207	14.8	1.01	76.3	1.97	205	70.86	13.5	0.80
	2	267	14.6	0.98	75.9	2.08	198	74.43	12.4	0.83
	3	0-100	-	-	-	-	-	-	-	-
	4	413	14.8	1.00	75.8	2.26	215	66.53	11.8	0.78
	ukp	892	14.7	0.98	75.9	1.86	204	68.39	13.4	0.98
	uks	131	14.6	1.03	74.8	1.98	185	49.61	12.2	0.70
Eastern	1	5,619	14.9	0.96	76.2	2.17	271	66.22	13.4	0.89
	2	2,656	14.9	0.98	76.0	2.05	278	64.90	12.5	1.01
	3	1,184	14.9	1.00	75.1	1.89	213	56.15	12.0	0.82
	4	3,573	14.9	0.97	75.0	2.15	226	71.36	11.9	0.80
	ukp	4,890	14.8	0.96	75.9	2.12	278	63.00	13.0	1.07
	uks	2,233	14.9	0.99	75.1	1.94	213	55.98	11.9	0.82
Midlands	1	917	14.8	0.94	76.7	2.10	282	55.77	13.4	0.90
	2	350	15.0	0.94	76.5	2.01	296	56.83	12.6	0.99
	3	0-100	-	-	-	-	-	-	-	-
	4	514	14.9	0.92	76.0	2.19	262	69.69	12.0	0.76
	ukp	791	14.9	0.92	76.5	2.08	294	52.41	13.2	1.05
	uks	225	14.8	0.90	75.2	1.96	234	50.32	12.1	0.82
North	1	453	14.8	1.00	77.3	2.04	280	50.60	12.8	0.86
	2	145	14.9	1.13	76.6	2.13	282	55.07	11.9	0.92
	3	0-100	-	-	-	-	-	-	-	-
	4	350	14.9	1.05	76.1	2.16	261	57.93	11.5	0.68
	ukp	268	14.8	1.05	76.8	2.12	283	49.17	12.5	1.06
	uks	213	14.8	1.08	75.7	2.05	231	43.25	11.5	0.62
GB	1	11,426	14.9	1.00	76.3	2.16	254	75.25	13.4	0.87
	2	3,949	14.8	0.98	76.1	2.08	271	70.61	12.5	1.02
	3	1,848	14.9	1.00	75.2	1.98	212	59.41	12.0	0.79
	4	5,466	14.9	0.99	75.3	2.23	231	71.54	11.9	0.80
	ukp	9,213	14.8	0.98	76.0	2.12	260	72.90	13.2	1.05
	uks	3,584	14.8	1.00	75.2	2.02	213	57.30	11.9	0.82

* Standard Deviation

All averages are simple arithmetic means

Results shown for 100+ samples only

Analysis by Harvest Date

Mean Values

Region	Harvest Date	Moisture Content		Specific Weight		Hagberg Falling Number		Protein Content	
		% d.m	s.d*	Kg/hl	s.d*	Seconds	s.d*	% d.m	s.d*
Eastern	Late July	-	-	-	-	-	-	-	-
	Early Aug	15.1	0.96	76.0	2.15	267	64.25	12.7	1.01
	Late Aug	14.9	1.01	75.2	2.09	237	68.89	12.5	1.10
	Early Sep	14.8	0.95	75.4	2.09	244	69.60	12.5	1.07
	Late Sep	14.9	0.91	75.6	2.05	244	70.69	12.7	1.09
	Early Oct	15.1	0.91	75.7	2.16	239	68.14	12.5	1.14
Northern	Late July	-	-	-	-	-	-	-	-
	Early Aug	15.0	1.12	77.0	2.05	285	51.84	12.3	0.99
	Late Aug	14.9	1.07	76.8	2.12	266	53.65	12.2	0.97
	Early Sep	14.8	1.05	76.4	2.26	269	54.99	12.0	0.96
	Late Sep	14.7	0.99	76.4	2.22	259	62.48	12.0	1.00
	Early Oct	-	-	-	-	-	-	-	-
South Eastern	Late July	-	-	-	-	-	-	-	-
	Early Aug	14.6	1.01	77.8	2.40	328	49.72	13.8	0.96
	Late Aug	15.2	1.16	76.4	2.27	238	83.93	13.4	0.95
	Early Sep	14.9	1.01	76.0	2.13	227	79.93	13.1	1.04
	Late Sep	14.7	1.03	75.7	2.26	221	77.48	13.0	1.02
	Early Oct	14.6	1.00	76.2	1.98	223	80.09	12.9	1.10
South Western	Late July	-	-	-	-	-	-	-	-
	Early Aug	15.1	1.21	76.2	2.33	233	73.59	13.5	0.91
	Late Aug	14.9	0.96	76.3	1.97	199	67.90	13.0	1.05
	Early Sep	14.6	1.00	75.9	2.07	202	68.62	12.8	1.10
	Late Sep	14.5	0.88	76.1	1.95	193	63.35	12.8	1.03
	Early Oct	-	-	-	-	-	-	-	-
Midlands	Late July	-	-	-	-	-	-	-	-
	Early Aug	15.1	1.06	77.3	2.09	301	54.39	13.2	1.00
	Late Aug	14.9	0.93	76.3	2.16	275	61.54	12.9	1.07
	Early Sep	14.8	0.92	76.1	2.11	271	60.41	12.6	1.08
	Late Sep	14.7	0.77	76.0	2.13	266	60.27	12.4	0.93
	Early Oct	-	-	-	-	-	-	-	-

* Standard Deviation

All averages are simple arithmetic means

Results shown for 100+ samples only

Wheat Analysis by Region

	Number of Samples	Moisture Content		Specific Weight		Hagberg F-Number		Protein Content	
		% d.m	s.d*	Kg/hl	s.d*	Seconds	s.d*	% d.m	s.d*
		Mean Values							
Eastern	13,855	14.9	0.97	75.7	2.19	255	70.43	12.6	1.10
Midlands	2,049	14.9	0.93	76.4	2.17	277	61.19	12.8	1.07
Northern	1,103	14.9	1.05	76.6	2.18	268	55.44	12.1	0.99
S. East	4,996	14.8	1.05	76.0	2.21	230	81.12	13.1	1.04
S. West	1,996	14.7	1.01	76.1	2.05	205	70.18	13.0	1.07
Wales	0-100	-	-	-	-	-	-	-	-
Scotland	214	14.1	0.93	77.1	2.42	229	74.07	11.4	0.90
GB	24,252	14.9	1.00	75.9	2.20	248	73.59	12.8	1.10

* Standard Deviation

All averages are simple arithmetic means

Results shown for 100+ samples only

Analysis by Variety, by Region

Variety	Region	Number of Samples	Moisture Content		Specific Weight		Hagberg F-Number		Protein Content	
			% d.m	s.d*	Kg/hl	s.d*	Seconds	s.d*	% d.m	s.d*
			Mean Values							
Skyfall	Eastern	1,000+	15.0	0.97	76.6	2.07	258	69.65	13.1	0.89
	Midlands	250-499	14.9	0.97	77.2	1.91	267	60.06	13.1	0.91
	Northern	250-499	14.8	1.01	77.6	1.86	276	52.35	12.6	0.81
	S. East	1000+	14.9	1.10	76.9	2.08	228	89.30	13.4	0.84
	S. West	500-999	14.7	1.05	76.9	1.98	197	78.13	13.3	0.81
	Scot	0-99	-	-	-	-	-	-	-	-
	GB	1000+	14.9	1.01	76.8	2.06	248	76.01	13.2	0.89
Crusoe	Eastern	1000+	14.8	0.96	75.6	2.18	268	59.37	13.8	0.71
	Midlands	250-499	14.8	0.88	76.5	2.11	291	46.57	13.7	0.78
	Northern	100-249	14.7	0.99	77.1	2.07	280	39.85	13.1	0.81
	S. East	1000+	14.8	1.02	75.9	2.17	223	79.30	13.6	0.79
	S. West	500-999	14.8	0.96	75.8	1.75	203	64.29	13.8	0.70
	Scot	0-99	-	-	-	-	-	-	-	-
	GB	1000+	14.8	0.98	75.8	2.14	245	72.85	13.7	0.75
Cordiale	Eastern	500-999	14.8	0.97	77.0	1.95	307	51.44	13.3	0.76
	Midlands	100-249	15.1	0.93	77.8	1.98	327	49.30	13.3	0.81
	Northern	0-99	-	-	-	-	-	-	-	-
	S. East	100-249	14.8	0.98	77.7	1.96	285	82.61	13.8	0.74
	S. West	0-99	-	-	-	-	-	-	-	-
	Scot	0-99	-	-	-	-	-	-	-	-
	GB	1000+	14.8	0.97	77.2	1.96	304	59.05	13.4	0.79
KWS-Siskin	Eastern	500-999	14.8	0.97	75.5	1.96	247	70.66	12.2	0.85
	Midlands	100-249	15.0	0.98	76.1	1.67	265	56.04	12.2	0.72
	Northern	0-99	-	-	-	-	-	-	-	-
	S. East	100-249	14.8	0.96	76.2	2.02	228	72.15	12.0	0.84
	S. West	100-249	14.6	0.91	74.9	1.79	148	49.47	12.3	0.76
	Scot	0-99	-	-	-	-	-	-	-	-
	GB	1000+	14.8	0.97	75.6	1.97	240	72.85	12.2	0.83
KWS-Lili	Eastern	500-999	14.9	0.98	75.5	1.85	276	57.01	11.9	0.82
	Midlands	0-99	-	-	-	-	-	-	-	-
	Northern	0-99	-	-	-	-	-	-	-	-
	S. East	100-249	14.9	1.02	75.1	1.74	214	69.20	12.2	0.77
	S. West	0-99	-	-	-	-	-	-	-	-
	Scot	0-99	-	-	-	-	-	-	-	-
	GB	1000+	14.9	1.01	75.5	1.85	265	64.66	12.0	0.81

* Standard Deviation

All averages are simple arithmetic means

Results shown for 100+ samples only

While the Agriculture and Horticulture Development Board seeks to ensure that the information contained within this document is accurate at the time of printing, no warranty is given in respect thereof and, to the maximum extent permitted by law the Agriculture and Horticulture Development Board accepts no liability for loss, damage or injury howsoever caused (including that caused by negligence) or suffered directly or indirectly in relation to information and opinions contained in or omitted from this document. AHDB Cereals and Oilseeds is part of the Agriculture and Horticulture Development Board.
© Agriculture and Horticulture Development Board 2017. All rights reserved.

AHDB T +44 (0) 247 647 8730
Stoneleigh Pe F +44 (0) 247 647 8904
Kenilworth cereals.mi@ahdb.org.uk
CV8 2TL cereals.ahdb.org.uk
UK



CEREAL QUALITY SURVEY RESULTS 2017

Barley Analysis by Variety

Mean Values

	Number of Samples	Moisture Content		Specific Weight		Nitrogen Content		Screening Values % through 2.25mm		Screening Values % retained 2.5mm	
		% d.m	s.d*	Kg/hl	s.d*	% d.m	s.d*	% d.m	s.d*	% d.m	s.d*
Winter	4,590	15.3	1.66	65.3	2.08	1.68	0.18	2.7	1.46	91.0	4.44
SY-Venture	871	14.9	1.48	65.3	1.95	1.68	0.14	3.4	1.43	89.4	3.93
Pearl	871	16.4	1.76	65.9	1.70	1.73	0.17	2.0	1.04	93.7	3.10
Maris Otter	663	14.6	1.27	66.5	1.99	1.55	0.15	3.6	1.32	87.4	3.60
Spring	23,025	16.1	1.85	64.2	1.99	1.68	0.20	1.7	1.10	94.7	3.03
Concerto	8,732	17.1	1.89	64.3	1.97	1.52	0.12	2.0	1.03	94.8	2.57
Propino	7,445	15.2	1.34	64.6	1.92	1.84	0.13	1.3	0.96	95.5	3.06
RGT-Planet	2,768	15.4	1.41	63.7	1.89	1.79	0.13	2.0	1.23	93.1	3.74

* Standard Deviation

*All averages are simple arithmetic means
Results shown for 100+ samples only*

Barley Analysis by Region

Mean Values

	Number of Samples	Moisture Content		Specific Weight		Nitrogen Content		Screening Values % through 2.25mm		Screening Values % retained 2.5mm	
		% d.m	s.d*	Kg/hl	s.d*	% d.m	s.d*	% d.m	s.d*	% d.m	s.d*
Eastern	7,017	15.0	1.18	64.2	2.03	1.77	0.17	1.9	1.36	93.6	4.23
Midlands	1,161	15.2	1.38	64.7	2.06	1.87	0.13	1.8	1.16	94.2	3.70
Northern	2,120	16.0	1.59	64.7	2.04	1.71	0.17	2.1	1.19	93.4	3.56
S. East	3,441	15.2	1.45	64.5	1.95	1.82	0.13	1.4	1.14	94.9	3.51
S. West	2,430	15.2	1.59	64.6	2.14	1.79	0.13	1.7	1.29	94.0	3.97
Wales	0-100	-	-	-	-	-	-	-	-	-	-
Scot	11,425	17.1	1.85	64.4	2.05	1.53	0.13	2.0	1.07	94.6	2.77
GB	27,615	15.9	1.84	64.4	2.04	1.68	0.19	1.9	1.23	94.2	3.49

* Standard Deviation

*All averages are simple arithmetic means
Results shown for 100+ samples only*

Analysis by Harvest Date - Barley

Mean Values

Region	Harvest Date	Moisture Content		Specific Weight		Nitrogen Content		Screening Values % through 2.25mm		Screening Values % retained 2.5mm	
		% d.m	s.d*	Kg/hl	s.d*	% d.m	s.d*	% d.m	s.d*	% d.m	s.d*
Eastern	Late July	14.7	1.07	65.5	2.06	1.61	0.17	3.0	1.46	89.5	4.14
	Early Aug	15.2	1.06	64.1	2.04	1.75	0.17	2.1	1.49	92.7	4.58
	Late Aug	15.0	1.18	64.0	1.78	1.83	0.14	1.6	1.09	94.8	3.42
	Early Sep	15.0	1.21	64.2	2.03	1.82	0.15	1.6	1.12	94.8	3.59
	Late Sep	15.2	1.22	63.8	1.94	1.82	0.13	1.6	1.16	94.8	3.32
	Early Oct	15.5	1.48	63.8	1.95	1.84	0.16	1.8	1.16	94.2	3.50
Northern	Late July	15.2	1.66	65.8	1.95	1.72	0.17	2.8	1.23	90.5	3.69
	Early Aug	16.3	1.57	64.4	1.82	1.76	0.16	2.2	1.21	92.6	3.65
	Late Aug	16.1	1.39	64.5	2.05	1.67	0.16	1.8	1.02	94.5	2.97
	Early Sep	16.1	1.60	64.4	1.96	1.71	0.17	2.1	1.17	94.2	3.04
	Late Sep	16.1	1.64	64.5	2.09	1.74	0.16	2.3	1.27	93.7	3.40
	Early Oct	-	-	-	-	-	-	-	-	-	-
Scotland	Late July	15.2	1.76	66.5	1.72	1.60	0.15	2.0	1.15	93.5	3.37
	Early Aug	16.5	1.98	64.8	1.91	1.63	0.18	1.6	1.02	94.5	3.11
	Late Aug	17.5	1.81	64.3	1.89	1.50	0.11	2.0	1.16	94.7	2.91
	Early Sep	17.2	1.69	64.3	1.95	1.52	0.11	2.1	1.06	94.5	2.55
	Late Sep	17.0	1.86	63.8	2.21	1.55	0.12	2.0	0.93	95.0	2.44
	Early Oct	17.1	2.10	62.7	2.31	1.56	0.12	2.2	1.00	94.8	2.91
South Eastern	Late July	14.0	1.49	66.0	2.12	1.67	0.15	3.4	1.43	88.5	4.04
	Early Aug	15.7	1.68	64.3	2.01	1.79	0.15	1.6	1.26	94.3	3.92
	Late Aug	15.3	1.40	64.5	1.81	1.82	0.12	1.2	0.95	95.3	3.03
	Early Sep	15.1	1.33	64.2	1.96	1.86	0.12	1.4	1.12	95.1	3.38
	Late Sep	15.0	1.12	64.6	1.82	1.84	0.12	1.4	0.90	95.1	3.12
	Early Oct	15.2	1.21	64.7	2.05	1.83	0.11	1.3	0.91	95.3	2.85
South Western	Late July	14.3	1.95	65.2	2.70	1.68	0.17	3.7	1.34	87.3	3.94
	Early Aug	15.5	1.74	64.5	2.06	1.79	0.13	1.9	1.43	93.2	4.19
	Late Aug	15.3	1.61	64.6	2.07	1.79	0.12	1.6	1.18	94.4	3.85
	Early Sep	15.1	1.38	64.4	2.14	1.81	0.12	1.5	1.07	94.7	3.41
	Late Sep	15.1	1.23	64.5	2.16	1.81	0.12	1.6	1.08	94.5	3.09
	Early Oct	14.9	1.00	64.7	2.14	1.82	0.12	1.3	0.89	94.7	2.93
Midlands	Late July	-	-	-	-	-	-	-	-	-	-
	Early Aug	15.7	1.34	64.0	1.93	1.89	0.12	2.1	1.26	92.4	3.96
	Late Aug	15.4	1.30	64.7	2.05	1.89	0.11	1.7	1.07	94.0	3.58
	Early Sep	15.1	1.25	64.8	2.02	1.90	0.11	1.6	1.01	95.0	3.03
	Late Sep	15.0	1.42	64.9	2.08	1.89	0.10	1.7	1.18	94.5	3.68
	Early Oct	-	-	-	-	-	-	-	-	-	-

* Standard Deviation

**All averages are simple arithmetic means
Results shown for 100+ samples only**

Analysis by Variety, by Region

Mean Values

Variety	Region	Number of Samples	Moisture Content		Specific Weight		Nitrogen Content		Screening Values % through 2.25mm		Screening Values % retained 2.5mm	
			% d.m	s.d*	Kg/hl	s.d*	% d.m	s.d*	% d.m	s.d*	% d.m	s.d*
SY-Venture	Eastern	250-499	14.6	1.08	65.2	1.90	1.66	0.14	3.4	1.44	88.5	3.69
	Midlands	0-99	-	-	-	-	-	-	-	-	-	-
	Northern	100-249	14.8	1.51	65.0	1.94	1.63	0.16	3.5	1.43	88.0	3.87
	S. East	0-99	-	-	-	-	-	-	-	-	-	-
	S. West	0-99	-	-	-	-	-	-	-	-	-	-
	Scot	100-249	16.6	1.35	65.9	1.19	1.71	0.12	0.0	0.00	92.2	2.96
	GB	500-999	14.9	1.48	65.3	1.95	1.68	0.14	3.4	1.43	89.4	3.93
Concerto	Eastern	500-999	15.0	1.18	63.4	2.05	1.66	0.16	2.4	1.48	92.2	4.22
	Midlands	0-99	-	-	-	-	-	-	-	-	-	-
	Northern	500-999	16.5	1.55	65.0	1.68	1.62	0.12	1.8	0.94	95.0	2.39
	S. East	0-99	-	-	-	-	-	-	-	-	-	-
	S. West	0-99	-	-	-	-	-	-	-	-	-	-
	Scot	1,000+	17.4	1.83	64.3	1.96	1.51	0.10	1.9	0.97	94.9	2.37
GB	1,000+	17.1	1.89	64.3	1.97	1.52	0.12	2.0	1.03	94.8	2.57	
Propino	Eastern	1,000+	15.0	1.16	64.4	1.81	1.83	0.14	1.3	0.98	95.4	3.22
	Midlands	500-999	15.3	1.30	65.1	1.92	1.90	0.10	1.3	0.89	95.5	2.89
	Northern	250-499	15.8	1.45	64.6	2.12	1.82	0.14	1.4	0.99	95.3	2.74
	S. East	1,000+	15.3	1.40	64.8	1.85	1.85	0.11	1.1	0.87	96.0	2.72
	S. West	1,000+	15.3	1.51	64.8	2.08	1.83	0.12	1.3	1.01	95.2	3.19
	Scot	0-99	-	-	-	-	-	-	-	-	-	-
	GB	1,000+	15.2	1.34	64.6	1.92	1.84	0.13	1.3	0.96	95.5	3.06
RGT-Planet	Eastern	1,000+	15.2	1.22	63.5	1.88	1.79	0.15	2.1	1.26	93.1	3.86
	Midlands	100-249	15.2	1.39	64.3	1.93	1.88	0.10	1.8	1.05	93.8	3.41
	Northern	100-249	15.5	1.46	64.1	1.70	1.75	0.14	2.2	1.22	93.2	3.31
	S. East	500-999	15.4	1.48	63.7	1.90	1.78	0.12	1.7	1.11	93.8	3.44
	S. West	250-499	15.5	1.68	63.7	1.85	1.75	0.11	2.2	1.32	91.9	3.93
	Scot	0-99	-	-	-	-	-	-	-	-	-	-
	GB	1,000+	15.4	1.41	63.7	1.89	1.79	0.13	2.0	1.23	93.1	3.74

* Standard Deviation

All averages are simple arithmetic means
Results shown for 100+ samples only

Analysis by Region, Winter/Spring

Region	Number of Samples	Moisture Content		Specific Weight		Nitrogen Content		Screening Values % through 2.25mm		Screening Values % retained 2.5mm	
		% d.m	s.d*	Kg/hl	s.d*	% d.m	s.d*	% d.m	s.d*	% d.m	s.d*
Eastern	7,017	15.0	1.18	64.2	2.03	1.77	0.17	1.9	1.36	93.6	4.23
Winter	1,609	14.7	1.06	65.4	2.07	1.65	0.18	3.1	1.45	89.3	4.12
Spring	5,408	15.1	1.19	63.9	1.92	1.80	0.15	1.7	1.19	94.4	3.71
Midlands	1,161	15.2	1.38	64.7	2.06	1.87	0.13	1.8	1.16	94.2	3.70
Winter	186	14.6	1.53	64.5	2.19	1.79	0.19	2.9	1.35	89.8	4.19
Spring	975	15.3	1.32	64.7	2.04	1.89	0.11	1.6	1.04	94.7	3.27
Northern	2,120	16.0	1.59	64.7	2.04	1.71	0.17	2.1	1.19	93.4	3.56
Winter	841	15.8	1.70	65.0	2.03	1.75	0.17	2.6	1.24	91.5	3.77
Spring	1,279	16.1	1.50	64.4	2.00	1.69	0.16	1.9	1.09	94.4	2.99
S. East	3,441	15.2	1.45	64.5	1.95	1.82	0.13	1.4	1.14	94.9	3.51
Winter	281	14.3	1.40	65.5	2.28	1.72	0.18	3.3	1.40	88.8	3.86
Spring	3,160	15.3	1.43	64.4	1.90	1.83	0.12	1.3	0.99	95.2	3.12
S. West	2,430	15.2	1.59	64.6	2.14	1.79	0.13	1.7	1.29	94.0	3.97
Winter	242	14.3	1.64	65.6	2.70	1.69	0.17	3.7	1.39	87.2	3.77
Spring	2,188	15.3	1.54	64.5	2.06	1.80	0.12	1.6	1.16	94.4	3.64
Scot	11,425	17.1	1.85	64.4	2.05	1.53	0.13	2.0	1.07	94.6	2.77
Winter	1,427	16.3	1.67	65.5	1.93	1.65	0.16	2.3	1.45	92.7	4.32
Spring	9,998	17.3	1.85	64.2	2.00	1.52	0.11	2.0	0.98	94.8	2.38
GB	27,615	15.9	1.84	64.4	2.04	1.68	0.19	1.9	1.23	94.2	3.49
Winter	4,590	15.3	1.66	65.3	2.08	1.68	0.18	2.7	1.46	91.0	4.44
Spring	23,025	16.1	1.85	64.2	1.99	1.68	0.20	1.7	1.10	94.7	3.03

* Standard Deviation

**All averages are simple arithmetic means
Results shown for 100+ samples only**

While the Agriculture and Horticulture Development Board seeks to ensure that the information contained within this document is accurate at the time of printing, no warranty is given in respect thereof and, to the maximum extent permitted by law the Agriculture and Horticulture Development Board accepts no liability for loss, damage or injury howsoever caused (including that caused by negligence) or suffered directly or indirectly in relation to information and opinions contained in or omitted from this document. AHDB Cereals and Oilseeds is part of the Agriculture and Horticulture Development Board.

© Agriculture and Horticulture Development Board 2017. All rights reserved.

AHDB
Stoneleigh Park
Kenilworth
CV8 2TL
UK

T +44 (0) 247 647 8730
F +44 (0) 247 647 8904
cereals.mi@ahdb.org.uk
cereals.ahdb.org.uk